

**Programmatic Arthropod Monitoring at
the Haleakalā High Altitude Observatories
and Haleakalā National Park and
Annual Inspection of the DKIST Facilities
and Grounds, Maui, Hawai'i**

September 2017

Prepared for

**K.C. Environmental, Inc.
P.O. Box 1208
Makawao, HI 96768
(808) 573-1903**



Pacific Analytics, L.L.C.

**35891 Richardson Gap Road
Scio, Oregon 97374
(541) 258-5919
*www.statpros.com***

Prepared by:

Pacific Analytics, L.L.C.
35891 Richardson Gap Road
Scio, Oregon 97374
Tel. (541) 258-5919
mail@statpros.com
www.statpros.com

Gregory Brenner
Senior Associate / Project Manager

William Haines
Associate Scientist

The pictures contained in this report are for the exclusive use by Pacific Analytics, L.L.C and its clients. All photographs are copyrighted by Pacific Analytics, L.L.C. and may not be reproduced or used without the express written permission of Pacific Analytics, L.L.C.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories and Haleakalā National Park Maui, Hawai‘i

I. TABLE OF CONTENTS

	Page
I. TABLE OF CONTENTS	1
II. EXECUTIVE SUMMARY	2
III. INTRODUCTION	4
IV. QUESTIONS OF INTEREST	9
V. METHODS	19
VI. RESULTS AND DISCUSSION	28
VII. LITERATURE CITED	53
APPENDIX A HO ARTHROPOD SPECIES LIST	59
APPENDIX B DKIST ARTHROPOD SPECIES LIST	62
APPENDIX B HALE ES ARTHROPOD SPECIES LIST	64

II. EXECUTIVE SUMMARY

The National Science Foundation (NSF) has authorized the development of the Daniel K. Inouye Solar Telescope (DKIST), previously known as the Advanced Technology Solar Telescope (ATST)) within the 18-acre University of Hawai‘i Institute for Astronomy High Altitude Observatories (HO) site. The DKIST represents a collaboration of 22 institutions, reflecting a broad segment of the solar physics community. The DKIST project will be the largest and most capable solar telescope in the world. It will be an indispensable tool for exploring and understanding physical processes on the Sun that ultimately affect Earth. The DKIST Project will be contained within a 0.74 acre site footprint in the HO site. An Environmental Impact Statement was completed for the DKIST project (NSF 2009), and the NSF issued a Record of Decision in December of 2009.

The Haleakalā National Park (HALE) Road Corridor is being used for transportation during construction and use of the DKIST. The HO and HALE road corridor contain biological ecosystems that are both unique and fragile. The landscape at HO is considered to be an alpine dry shrubland vegetation type and resources along the Park road corridor are

grouped into alpine and subalpine shrubland habitat zones, depending upon the elevation. These habitats contain several native and non-native species of plants, animals, and arthropods. While the overall impacts on Hawaiian native arthropod resources within the Park road corridor during the construction phase would be considered minor, NSF has committed to several mitigation measures to reduce the impacts to these biological resources, including programmatic monitoring for active preservation of invertebrates before, during and after construction of the DKIST Project.

After preliminary sampling near the HALE Entrance Station and at the DKIST site in 2009, Programmatic Arthropod Monitoring and Assessment at the Haleakalā High Altitude Observatories and Haleakalā National Park was initiated with two sampling sessions in 2010. Monitoring is being conducted twice a year during the construction phase of the DKIST which began in December 2012. Semi-annual monitoring has occurred in 2011, 2012, 2013, 2014, 2015, 2016 and 2017.

This report presents the results of the Summer 2017 sampling. The goal is to

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

monitor the arthropod fauna at the DKIST site and along the HALE Road Corridor, identify Hawaiian native arthropod species or habitats, if any, that may be impacted by construction of the DKIST, and detect and identify alien invasive arthropod species that could have adverse impacts on the flora and fauna on Haleakalā. Programmatic Arthropod Monitoring studies are being coordinated and conducted with the approval of HALE.

This monitoring project provides a means of gathering reliable information that can be used to protect the native Arthropod species during development of observatory facilities and supports astronomy programs at the Haleakala High Altitude Observatory Site by promoting the good stewardship of the natural resources located there.

In addition to semi-annual monitoring required by the FEIS, pursuant to the approved HCP and published BO, an inspection for non-indigenous arthropod species is required to be conducted on an ongoing annual basis during the

construction phase and 50 year lifespan of the DKIST for programmatic monitoring. Facilities and grounds within 100 feet of the DKIST observatory buildings are to be thoroughly inspected for introduced species that may have eluded the cargo inspection processes or transported to the site by construction personnel.

This report also describes the results of the Annual Inspection conducted in September 2017. The goal is to verify that the DKIST is in compliance with the conditions and mitigation measures described in the guiding environmental documents.

The 2017 Annual Inspection was conducted on September 7, 2017. No non-indigenous, invasive arthropods were found at the site or on any of the construction material and equipment. The project was found to be compliant with all the mitigation measures in the guiding environmental documents for construction of the DKIST. The construction site and surrounding lay-down/storage areas were clean and free of non-indigenous invasive arthropods.

III. INTRODUCTION

Programmatic Monitoring

The Haleakalā volcano on the island of Maui is one of the highest mountains in Hawai‘i, reaching an elevation of 10,023 feet (3,055 m) at its summit on Pu‘u ‘Ula‘ula. Near the summit is a volcanic cone known as Kolekole with some of the best astronomy viewing in the world.

The National Science Foundation (NSF) has authorized the development of the Daniel K. Inouye Solar Telescope (DKIST), previously known as the Advanced Technology Solar Telescope (ATST)) within the 18-acre University of Hawai‘i Institute for Astronomy High Altitude Observatories (HO) site. The DKIST represents a collaboration of 22 institutions, reflecting a broad segment of the solar physics community. The DKIST project will be the largest and most capable solar telescope in the world. It will be an indispensable tool for exploring and understanding physical processes on the Sun that ultimately affect Earth.

The DKIST Project will be contained within a 0.74 acre site in the HO site. An Environmental Impact Statement was completed for the DKIST project (NSF 2009), and the NSF issued a Record of Decision in December of 2009. The

Haleakalā National Park (HALE) Road Corridor is being used for transportation during construction and use of the DKIST. Construction began in December 2012 and was ongoing during the Summer 2017 sampling.

The HO and HALE road corridor contain biological ecosystems that are both unique and fragile. The landscape at HO is considered to be an alpine dry shrubland vegetation type. A diverse fauna of resident insects and spiders reside there (Medeiros and Loope 1994). Some of these arthropods inhabit unique natural habitats on the bare lava flows and cinder cones with limited vegetation. Vegetation covers less than 5% of the open ground, and food is apparently scarce.

The ecosystem at the HO is extremely xeric, caused by relatively low precipitation, porous lava substrates that retain negligible amounts of moisture, little plant cover, and high solar radiation. The dark, heat-absorbing cinder provides only slight protection from the extreme temperatures. Thermal regulation and moisture conservation are critical adaptations of arthropods that occur in this unusual habitat.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

An inventory and assessment of the arthropod fauna at the HO site was conducted in 2003 as part of the Long Range Development Plan (LRDP) for the Haleakalā High Altitude Observatories. This inventory and assessment was updated in December 2005 to provide a more detailed description of the arthropod fauna at the two proposed DKIST sites, and identify Hawaiian native arthropod species or habitats, if any, that could be impacted by construction of the DKIST. In an effort to be complete, supplemental sampling was conducted in 2007 to provide a seasonal component and additional nighttime sampling not included in the previous two inventories. Sampling in June 2009 was conducted to establish baseline conditions for future Programmatic Monitoring.

The landscape along the HALE road corridor is classified as alpine and subalpine shrubland habitat zones, depending upon the elevation. These habitats contain several native and non-native species of plants, animals, and arthropods. The subalpine shrubland within the Haleakalā National Park is also host to a wide variety of indigenous arthropod species (Krushelnycky et al. 2007). The vegetation there covers most of the open ground, mostly with native trees and shrubs, with native and alien grasses growing between. Precipitation in the

form of rainfall and fog is frequent, with about 70 inches falling throughout the year (Giambelluca et al. 1986).

While the overall impacts on arthropod resources within the Park road corridor during the construction phase would be considered minor, NSF has committed to several mitigation measures to reduce the impacts to these biological resources, including programmatic monitoring for active preservation of invertebrates during and after construction of the DKIST Project.

Environmental monitoring is the scientific investigation of the changes in environmental phenomena, attributes and characteristics that happen over time. Ecosystems are dynamic. Habitat conditions change daily, seasonally, and over longer periods of time. Animal and plant populations rise or fall in response to a host of environmental fluctuations. The general purpose of monitoring is to detect, understand, and predict the biological changes.

The scientific scope of the current phase of Arthropod Monitoring is to repeatedly sample arthropod habitats that may be impacted by construction of the DKIST, document changes to native arthropod populations, and detect new or potentially threatening invasive species of

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

arthropods that may impact the native resident arthropod fauna. Programmatic Arthropod Monitoring includes identification and taxonomy for both ground and shrub dwellers and is being conducted in both developed and undeveloped areas of HO (excluding the Air Force site).

Arthropod Programmatic Monitoring consists of one week sampling sessions conducted in the Summer and Winter months using standard arthropod sampling methods similar to those used during the 2007 inventory of arthropods within HALE (Kruschelnycky et al. 2007), collecting invertebrates both day and night, with identification and taxonomy for both ground and shrub dwellers in developed and undeveloped portions of the sampling areas.

The primary areas being sampled are the Haleakalā High Altitude Observatories (HO) site on Kolekole Hill, but not including the Air Force site, the DKIST Construction Site, and selected portions of the HALE Road Corridor. The 18 acre HO facility hosts several existing observatories and their support buildings, and also includes several undeveloped sites where native vegetation and the associated arthropod fauna is relatively undisturbed. DKIST construction is currently taking place in

previously undisturbed land located east of the existing Mees Solar Observatory facility. The portions of the HALE Road Corridor being sampled are determined in collaboration with the HALE staff biologists at the beginning of each sampling session.

Programmatic Monitoring will provide much of the data needed to protect and enhance natural resources, to modify management actions, to aid in compliance with environmental statutes, and to enhance public education and appreciation of the natural resources at the summit of Haleakalā.

The nomenclature used in this report follows the Hawaiian Terrestrial Arthropod Checklist, Third Edition (Nishida 1997) and the Manual of the Flowering Plants of Hawai‘i (Wagner and others 1990). Hawaiian and scientific names are italicized unless major taxonomic revisions were available.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Species are discussed as being endemic, indigenous, non-indigenous, adventive, and purposely introduced. These terms are defined as:

Endemic – A species native to, or restricted to Hawai‘i.

Indigenous – A species native to Hawai‘i but that naturally occurs outside of Hawai‘i as well.

Non-indigenous – A species not native to Hawai‘i.

Adventive – Not native, a species transported into a new habitat by natural means or accidentally by human activity.

Purposely introduced – A species released in Hawai‘i for a particular purpose, usually to control a weedy plant or another insect.

This report describes the results of sampling conducted in September 2017, the second of two sampling sessions for Programmatic Arthropod Monitoring and Assessment this year, and continues monitoring that began in September 2009. The goal is to monitor the arthropod fauna at the HO site, the DKIST construction site, and along the selected portions of the HALE Road Corridor, identify Hawaiian native arthropod species or habitats, if any, that may be impacted by construction of the DKIST, and detect and identify alien invasive arthropod species that could have adverse

impacts on the flora and fauna on Haleakalā. Programmatic Arthropod Monitoring studies are being coordinated and conducted with the approval of HALE staff biologists.

Sampling of arthropod habitats was approved in a permit obtained from the Department of Land and Natural Resources (Endorsement No. I1014), effective date February 1, 2017 – February 1, 2018, and the National Park Service (Permit # HALE-2015-SCI-0003) issued on April 16, 2015. Sampling began on September 2, 2017 and was completed on September 10, 2017.

Annual Inspection

An inspection is required to be conducted on an ongoing annual basis during the construction phase and 50 year lifespan of the DKIST. The inspection was conducted on September 7, 2017. DKIST interior facilities and grounds within 100 feet of the buildings are to be thoroughly inspected for introduced species that may have eluded the cargo inspection processes or transported to the site by construction personnel. Any newly-discovered non-native, invasive arthropod are to be photo documented, mapped, and described. Arrangements will be made for eradication of any invasive introduced species found inside or within 100 feet of the DKIST buildings.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Appropriate control methods include the use of available herbicides and pesticides, in accordance with established practice at HO (University of Hawai‘i 2010) and pursuant to label requirements.

Annual Inspections provide information about compliance with the guiding environmental documents prepared for

the DKIST project. These documents include the DKIST Habitat Conservation Plan, USFWS biological Opinion, and the DKIST FEIS. In addition, the inspection meets the requirements of the University of Hawai‘i Institute for Astronomy Management Plan, which describes mitigation measures to prevent introduction of introduced species.



Native Arthropod Habitat adjacent to the DKIST construction lay-down area.
Photo taken September 7, 2017.

IV. QUESTIONS OF INTEREST

Important Questions of Interest are those with answers that can be efficiently estimated and that yield the information necessary for management decision-making. The following Questions of Interest were developed for Programmatic Monitoring and the Annual Inspection, and are the focus of this report.

Programmatic Monitoring

Question 1

What are the characteristic arthropod populations at the DKIST site, the larger HO site (excluding the Air Force site), and along selected areas of the HALE Road Corridor?

Justification:

Programmatic Monitoring will yield a comprehensive list of the characteristic arthropod fauna at the DKIST site, developed and undeveloped areas of the HO site, and along selected areas of the HALE Road Corridor.

Monitoring goals:

- 1) To describe the characteristic arthropod populations at the DKIST site, the larger HO site, and along the HALE Road Corridor,
- 2) To provide historical records of change in native arthropod species population attributes, and characteristics.

The results of this sampling are combined with information gathered during previous studies to develop a comprehensive list of arthropods at the Haleakalā High Altitude Observatories (HO) site, the DKIST site, and along selected areas of the HALE Road Corridor, and a qualitative description of seasonal variations in their abundance.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Question 2

What adverse impacts can be detected, if any, on characteristic populations of arthropods at the DKIST site, the larger HO site (excluding the Air Force site), and along selected areas of the HALE Road Corridor that may be due to DKIST construction?

Justification:

Programmatic Monitoring of native arthropod species will yield reliable scientific information about the current status (presence and abundance) of these species at the sampling sites. The information will be useful to detect changes and trends that may be due to the construction of the DKIST.

Monitoring goals:

- 1) To detect changes, trends, periodicities, cycles, and/or other patterns of change in arthropods at the DKIST site, the larger HO site, and along the HALE Road Corridor during the construction of the DKIST.

Programmatic Monitoring reports provide a discussion of the results of sampling, a description of changes in presence or abundance, and an assessment of those changes that may be due to the DKIST construction, and provide opportunities for adaptive management of construction processes, through the use of control measures, where these changes and/or trends negatively affect the arthropod population.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Question 3

What non-indigenous invasive arthropod species, if any, are detected at the DKIST site, the larger HO site (excluding the Air Force site), and along selected areas of the HALE Road Corridor during DKIST construction?

Justification:

Programmatic Monitoring for non-indigenous invasive arthropod species will detect potential threats to the nearby native ecosystems before they have an opportunity to establish resident populations. Early detection will allow implementation of control measures to eradicate invasive arthropod species (e.g. ants and spiders) before they can damage the nearby native ecosystems.

Monitoring goals:

- 1) To detect non-indigenous invasive arthropod species at the DKIST site, the larger HO site, and along selected areas of the HALE Road Corridor during construction of the DKIST.

If any invasive arthropod species (e.g. ants and spiders) are detected, eradication measures will be implemented to prevent these species from establishing resident populations.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Annual Inspection

Question 4

What non-indigenous arthropod species, if any, are detected within the interior of DKIST facilities and the grounds within 100 ft. (30 m) of the buildings?

Justification:

Detailed site inspections for non-indigenous arthropod species can detect potential threats to the nearby native ecosystems that may have escaped detection during the regular inspection process or programmatic monitoring. Early detection will allow implementation of control measures to eradicate invasive arthropod species (e.g. ants and spiders) before they have an opportunity to establish resident populations that could damage nearby native ecosystems.

Inspection goals:

- 1) To detect invasive non-indigenous arthropod species inside of the DKIST facilities and the grounds within 100 feet of the buildings.

If any invasive arthropod species (e.g. ants and spiders) are detected, they will be photo documented, mapped, and described, and then exterminated. Eradication measures may include brushing away spider webs to disrupt mating and foraging, sticky traps to capture ants, and the application of pesticides in accordance with established practice at HO (University of Hawai‘i 2010) and pursuant to label requirements.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Question 5

What non-indigenous invasive arthropod species, if any, are detected at the DKIST construction, lay-down and staging areas?

Justification:

While these areas are sampled during Programmatic Monitoring, they will receive increased scrutiny during the Annual Inspection. Annual inspection for non-indigenous arthropod species will detect potential threats to the nearby native ecosystems that may have escaped earlier detection and before they have an opportunity to establish resident populations. Early detection will allow implementation of control measures to eradicate invasive arthropod species (e.g. ants and spiders) before they can damage the nearby native ecosystems.

Inspection goals:

- 1) To detect invasive non-indigenous arthropod species at the DKIST lay-down and staging areas.

If any invasive arthropod species (e.g. ants and spiders) are detected, they will be photo documented, mapped, and described, and then eradicated to prevent these species from establishing resident populations. Eradication measures may include brushing away spider webs to disrupt mating and foraging, sticky traps to capture ants, and the application of pesticides in accordance with established practice at HO (University of Hawai‘i 2010) and pursuant to label requirements.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Question 6

Are mitigation measures implemented that prevent the establishment of invasive species due to DKIST construction activities?

Justification:

NSF has committed to several mitigation measures described in the DKIST FEIS, Habitat Conservation Plan (HCP), and USFWS Biological Opinion (BO) to prevent the introduction of invasive species to those areas surrounding the DKIST construction activities. The Annual Inspection will include examination of the DKIST Construction Site to ensure mitigation measures are being implemented correctly.

Monitoring goals:

- 1) To confirm that mitigation measures to prevent the establishment of invasive non-indigenous arthropod species committed to in the DKIST FEIS, HCP, and BO are being implemented correctly.

If any violations of the mitigation measures are detected, they will be photo documented, mapped, and described, and then reported to the Construction Site Manager, who will arrange for proper implementation of the measures to prevent invasive species from establishing resident populations.

**Specific Alien Arthropod Control Measures to be taken
(Habitat Conservation Plan Page 54 – 57 and Biological Opinion Page 20-24)**

Alien arthropods can arrive at the site by two general pathways. First, alien species already on Maui can spread to new locations. Second, alien species can arrive on the island with construction materials in or on shipping crates and containers. In order to block the first pathway, heavy equipment, trucks, and trailers will be pressure-washed before being moved to the DKIST construction site. The following specific alien arthropod control measures, adapted from those already required pursuant to the HO Management Plan will be implemented to further minimize the spread and establishment of alien insects. These six specific alien arthropod control measures are as follows:

- 1) Earthmoving equipment will be free of large deposits of soil, dirt and vegetation debris that could harbor alien arthropods.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

shipping inspections. Re-inspection prior to transport to the site will be completed to reduce the potential for undetected arthropods to reach the construction site. Arrangements will be made stipulating mandatory use of the Maui Alien Species Action Plan (ASAP) building for complete inspection of all possible items. This will prevent /or best allow for alien species interdiction on arriving materials.

- i. Inspect construction materials before entering the Park: Alien arthropods already resident in Hawai‘i are capable of hitchhiking on construction material such as bricks and blocks, plywood, dimension lumber, pipes, and other supplies. Precautions will be taken to ensure that alien arthropods are not introduced to the HO site.
- ii. Construction materials will be inspected before transport to the construction site.

If any alien arthropods are discovered, the infestation will be removed prior to transport. Infestations of ants can be removed using pressure-washing. Infestations of spiders can be removed using brooms, vacuum cleaners, or other similar methods. Pesticide use on materials to be transported to the site should be avoided.

- 3) Sanitary control of food and garbage will prevent access to food resources that could be used by invading ants and yellow jackets. Outdoor trash receptacles will be secured to the ground, have attached lids and plastic liners, and their contents will be collected frequently to reduce food availability for alien predators. Heavy, hinged lids will be used to prevent wind dispersal of garbage. Refuse will be collected on a regular basis to ensure containers do not become full or overflow. This could entail collection several times a week, particularly in eating areas and during periods of heavy use of the area. Containers will be regularly washed using steam or soap to reduce odors that attract ants. Plastic bag liners will be used in all garbage containers receiving food to contain leaking fluids.
- 4) Ensure construction waste and debris is secured to ensure it is not dispersed.
 - a. Construction activity may generate a considerable amount of waste debris. Typically construction debris is disposed of in “roll-off” containers that are periodically picked up and emptied at a landfill. Large “roll-off” containers can accommodate debris generated over several days of construction. Debris disposed of in these containers consists of wood, scrap insulation, packaging material, waste concrete, and various other construction wastes.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

- b. High winds at the site can disperse construction debris from the containers and disperse the material into adjacent arthropod habitat. Unsecured building materials and equipment at the site are also susceptible to wind dispersal. Construction trash and building material is not believed to significantly impact native arthropod species, but collection of the wind-blown material could potentially disturb their habitat (e.g., Howarth, et al., 1999).
- c. Construction trash containers will be tightly covered to prevent construction wastes from being dispersed by wind. This will be accomplished during construction of DKIST pursuant to the best management practices described in the HO Management Plan.

Covering containers will decrease the amount of construction debris that could be blown onto adjacent native arthropod habitat. “Roll off” containers can be equipped with tarps held securely with cables. Containers will be collected on a regular basis before they are completely full or overflowing. This could entail collection several times a week, particularly during periods of heavy use.

- 5) Invasive species detection and interdiction will be the responsibility of the resource biologist for DKIST and supporting avian biologist. Detection and interdiction will be conducted routinely by these personnel to ensure that new introductions are controlled.
 - a. A biological monitor will be employed during construction and programmatic arthropod sampling will be done in accordance with the schedule described within Section 2.3-DKIST Project Description of the HCP, and Section 3.0 of the BO. Monitoring for new alien arthropod introductions will be conducted during construction activities and any populations detected will be eradicated. Monitoring for alien populations is relatively easy and inexpensive to conduct. Baited traps have been shown to detect alien populations before they reach damaging proportions.
 - b. Ant eradication: Sticky traps designed to capture ants will be deployed immediately after any ants are detected. Persistence of ant detections are indicative of larger infestations, and will prompt a search for and eradication of colonies. Bait and chemical control will be employed only when absolutely necessary and only by a certified pest control professional.
 - c. Alien spider eradication: Any alien spider webs detected will be removed. Native lycosid wolf spiders do not make webs. Native sheet-web spiders make tiny webs under the cinder surface. Only alien spiders would make large spider

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

webs at HO. Sweeping such webs away with a broom disrupts alien spider food capture success and destroys egg masses. Follow-up measures will be developed and implemented to control alien spiders when they are detected.

- 6) Construction materials stored at the site will be covered with tarps, or anchored in place, and will not be susceptible to movement by wind. Securing materials will reduce the chances of debris being dispersed from the site into native arthropod habitat. Construction materials and supplies will be prevented from being blown into native arthropod habitat by covering them with heavy canvas tarps, using steel cables, attached to anchors that are driven into the ground. Construction materials at the site will be tied down or otherwise secured during high winds and at close of work each day. If construction materials and trash are blown into native arthropod habitat, they will be collected with a minimum of disturbance to the habitat.

V. METHODS

Site Description

The Haleakalā High Altitude Observatories (HO) site is located on Kolekole Hill. The highest point on the HO site is at 3,052-m (10,012-ft) above sea level. The 7.3-ha (18.1-ac) site was established in 1961, and the first telescope, the Mees Solar observatory was dedicated in 1964. The site now consists of five substantial telescope facilities, in addition to several smaller facilities.

The DKIST site is on undeveloped land located east of the existing Mees Solar Observatory facility at 3,042-m (9,980-ft) above sea level. Annual precipitation averages 1,349.2-mm (53.14-in), falling primarily as rain and mist during the winter months from November through April. Snow rarely falls at the site.

Haleakalā sampling locations were determined with guidance and cooperation from HALE personnel. During this session, sampling was conducted in the area near the HALE Entrance Station, at about 2,072 m (6,800 ft) on the western slope of Haleakalā.

Monitoring Procedures

The selection of a trapping technique used in a study was carefully considered. When the target species of the trapping system are rare or important for other reasons (i.e., endangered, keystone species, etc.) live-trapping should be considered. Entomologists have long believed that they can sample without an impact on the population being sampled. It has been assumed that collecting has only a small impact on the populations of interest. While this assumption remains to be tested, responsible entomologists consider appropriate trapping techniques to ensure survival of local populations of interest. The sampling methods that were used during this study are similar to those used during the 2007 arthropod inventory conducted on the western slope of Haleakalā and were reviewed by HALE natural resource staff and modified according to their comments.

Pitfall Trapping

After consultation with HALE natural resources staff, ten pitfall traps were installed near HALE Entrance Station site (five below the road and five above the road). Ten pitfall traps were installed at the Haleakalā High Altitude Observatories (HO) site in both developed and undeveloped areas, and

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

ten pitfall traps were deployed at the DKIST site. The traps (300 ml [10 oz], 80 mm diameter cups) were filled with soapy water solution as preservative. Concerns about endangered native birds precluded the use of ethylene glycol. The traps were spaced at least 2 m apart, and left open for seven days at the DKIST site and for seven days at the HALE site. It was decided that pitfall traps would not be baited around the rim with blended fish because they might attract birds. This is a trapping method similar to that used during an arthropod survey conducted in 2007 (Krushelnycky et al. 2007).

Care was taken to avoid archeological sites. These sites have cultural and historical significance and precautions were made to prevent their disturbance. Traps were not placed in or near these

sites. A map of significant historic and cultural sites within 50 feet of the road corridor was used to avoid such sites. Habitat was accessed with a minimum of disturbance to the habitat. Care was also taken to prevent creation of new trails or evidence of foot traffic.

Care was also taken to avoid disturbing nesting petrels and other wildlife species. The endangered petrels dig into cinder to make burrows for nesting. Efforts were made to avoid known burrows. Pitfall traps are placed below ground and covered with a heavy cap rock. This makes it very unlikely that petrels could access the traps.

All pitfall traps were installed on September 2, 2017 and their contents collected on September 10, 2017.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i



Locations of pitfall traps (blue circles) and light traps (yellow circles) at the HALE Entrance Station.

Light-Trapping

Sampling for nocturnal insects is vital to understanding the complete faunal presence. Some insects are only active and

moving around at night. Many insects have a nocturnal activity cycle to evade birds, and to locate certain food sources. Night collecting is important in

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

environments like dry locations where insects may choose this strategy to avoid desiccation.

Battery-powered ultraviolet light traps were operated near the HALE Entrance Station, at the HO site, and at the DKIST site. The traps consisted of a 3.5 gallon polypropylene bucket, a smooth surface funnel, a 22 watt Circline blacklight tube mounted on top of vanes under an

aluminum lid that directs light downwards. The effective range of the 22 watt lamp is less than 100 feet, and traps were always located more than 100 feet from the nearest petrel burrow. Light traps were run for seven nights at the DKIST site, HO site, and the HALE site. Light traps were set at each sampling site near sunset, and were allowed to run overnight or until batteries failed.



Locations of pitfall traps (blue circles) and light traps (yellow circles) at the HO and DKIST.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Other Light Sampling at Night

Night collecting can be aided by a UV light source. Small handheld ultraviolet blacklights were used for additional sampling for foliage and ground-dwelling arthropods.

Visual Observations and Habitat Collecting Under Rocks and in Leaf Litter

Time was spent sampling under rocks, in leaf litter, and on foliage to locate and collect arthropods at each sampling station. Hand picking, while sorting through leaf litter and bunch grasses, and searching beneath stones was the most effective sampling for litter and soil associated forms.



Leaf litter and plant debris were placed in a plastic tub and searched for arthropods.

Collecting on Foliage

Foliage of various common plant species was sampled by beating sheet. A one-meter square beating sheet or insect net was placed under the foliage being sampled and the branch hit sharply three times using a small plastic pipe. After the initial collection the foliage was beaten again to dislodge persistent individuals. Care was taken to avoid sensitive plants and to leave vegetation intact.

Nets

Aerial nets and sweep nets were used as necessary to capture flying insects and arthropods that occur on grasses.

Baited Traps

Baited traps were deployed to detect the presence of ants. These traps consisted of an index card, baited with tuna, honey, and peanut butter, and weighted down with a small rock. Traps were set near areas where ants could be introduced or where they may be foraging for food. Baited traps were deployed on the HO and DKIST sites on three different days. The traps were checked after forty-five minutes at which time the traps were removed. Baited traps were not left open overnight in order to avoid attracting unwanted pests.

Fifty baited ant traps were deployed on September 3, 2017 at the HO/DKIST sites.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Fourteen baited traps were deployed at the HALE ES site on September 4, 2017.

Inspection of construction lay-down and storage areas

Construction material and equipment in developed lay-down areas were visually inspected for invasive arthropod species and evidence of their presence. Specifically, these areas were inspected for the presence of ants, spiders, spider webs, and indications of the presence of other potentially invasive arthropod species.

Population Estimates

Although NSF committed to “population estimates for developed and undeveloped areas within HO, the DKIST Construction Site, and selected areas of the HALE road corridor” (NSF 2009), they are not possible with the approved sampling techniques. A consultation with the NPS determined that any data collected would be only a snapshot in time, reflective only of the sites sampled, and that the results are seasonal and could not be extrapolated beyond those limits. They also expressed an opinion that any “population estimates” would not be comparable over time and that accurate population estimates for arthropods are not possible with the sampling methods approved for use. In consultation with NPS staff biologists, it was decided that

sampling results would be presented as presence/absence, and that qualitative abundance estimates would be a suitable substitute for “population estimates” described in the FEIS (NSF 2009).

Sampling results in this report are presented as presence/absence, and, for selected species, qualitative abundance estimates are substituted for “population estimates” described in the FEIS (NSF 2009).

Relative abundance categories are:

- *infrequent* (individuals captured or observed < 10),
- *uncommon* (10 < individuals captured or observed < 25),
- *common* (25 < individuals captured or observed < 100), and
- *abundant* (100 < individuals captured or observed).

It should be noted that abundance designations are based exclusively on the capture or observation of specimens encountered at the sampling sites visited during each sampling session, and may be biased against certain species. For example, some ground dwelling species may be under-sampled because traps will not be baited and therefore not attractive to these species. Other species may be more or less abundant at other times of year than those sampled, or not

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

efficiently captured with the sampling methods used. These species may generally be more or less common than indicated from the results. The results presented in reports are only snapshots in time, reflective only of the sites sampled, and the results are seasonal and should not be extrapolated beyond those limits.

Collections

Arthropods that appear in traps were stored and later mounted for identification. Arthropods that are observed during hand collecting and netting were collected only as necessary to provide accurate identification and voucher specimens.

Curation

The contents of the traps were cleaned in 70% ethyl alcohol and placed in vials. The specimens were sorted into the morphospecies for identification. Hard-bodied species, such as beetles, moths, true bugs, flies, and wasps were mounted on pins, either by pinning the specimen or by gluing the specimens to paper points. Pinned specimens were placed into Schmidt boxes. Soft-bodied specimens, such as spiders and caterpillars were stored in vials filled with 70% ethyl alcohol.

Identification

Specimens were mounted and identified to the lowest taxonomic level possible within the time frame of the study. Many small flies and micro-Hymenoptera were sorted to morphospecies and will be sent to reliable experts for identification. Identification of arthropods is difficult, even for experts. More time needs to be allotted for this necessary task in all arthropod inventory projects. All specimen identifications are provisional until they can be confirmed by comparison to museum specimens or by group/taxon experts.

References for general identification of the specimens included *Fauna Hawaiiensis* (Sharp (ed) 1899-1913) and the 17 volumes of *Insects of Hawai‘i* (Zimmerman 1948a, 1948b, 1948c, 1948d, 1948e, 1957, 1958a, 1958b, 1978, Hardy 1960, 1964, 1965, 1981, Tentorio 1969, Hardy and Delfinado 1980, Christiansen and Bellinger 1992, Liebherr and Zimmerman 2000, and Daly and Magnacca 2003). Other publications that were useful for general identification included *The Insects and Other Invertebrates of Hawaiian Sugar Cane Fields* (Williams 1931), *Common Insects of Hawai‘i* (Fullaway and Krauss 1945), *Hawaiian Insects and Their Kin* (Howarth and Mull 1992), and *An Introduction to the*

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Study of Insects Sixth Edition (Borror, Triplehorn, and Johnson 1989).

For specific groups specialized keys were necessary. Most of these had to be obtained through a research library. Keys used to identify Heteroptera included those by Usinger (1936, 1942), Ashlock (1966), Beardsley (1966, 1977), Gagné (1997), Polhemus (2002, 2005, 2011, 2014), and Asquith (1994, 1997). Keys used to identify Hymenoptera included Cushman (1944), Watanabe (1958), Townes (1958), Beardsley (1961, 1969, 1976), Yoshimoto and Ishii (1965), and Yoshimoto (1965a, 1965b).

Species identification of those specimens identified to genus or species levels are unconfirmed and subject to change after comparison to specimens in museums.

In many cases changes in family and generic status and species synonymies caused species names to change from those in the keys. Species names used in this report are those listed in *Hawaiian Terrestrial Arthropod Checklist Third Edition* (Nishida 1997) unless a recent major taxonomic revision was available.

Schedule/Start and End Dates

Sampling was conducted over nine days and seven nights beginning on September 2, 2017 and ending on September 10, 2017.

Annual Inspection Procedures

Inspection of DKIST building interiors and within 100 feet of the DKIST buildings.

During the inspection, the interiors of all DKIST buildings were examined for evidence of non-indigenous invasive species. If any invasive arthropod species (e.g. ants and spiders) were detected, they would have been photo documented, mapped, and described, and then arrangements would be made for eradication to prevent these species from establishing resident populations. Eradication measures may include brushing away spider webs to disrupt mating and foraging, sticky traps to capture ants, and the application of pesticides in accordance with established practice at HO (University of Hawai‘i 2010) and pursuant to label requirements.

Inspection of DKIST construction, lay-down and staging areas.

During the Annual Inspection, construction lay-down and staging areas were examined for evidence of non-indigenous invasive species. If any invasive arthropod species (e.g. ants and spiders) were detected, they would have been photo documented, mapped, and described, and then arrangements would have been made for eradication to prevent these species from establishing resident populations. Eradication measures may include brushing away spider webs to disrupt mating and foraging, sticky traps to capture ants, and the application of pesticides in accordance with established practice at

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

HO (University of Hawai‘i 2010) and pursuant to label requirements.

Earthmoving Equipment

Earthmoving equipment and vehicles were inspected to verify they are being properly washed.

Construction materials, crates, shipping containers, packaging material, and observatory equipment

Construction materials, crates, shipping containers, packaging material, and observatory equipment were examined for evidence of non-indigenous arthropod species.

Sanitary control of food and garbage

Outdoor trash receptacles were examined to ensure they were secured to the ground, had attached lids and plastic liners. Containers were examined to verify that they were washed and that

odors were not present that may attract ants or yellow jackets.

Construction Waste and Debris

“Roll-off” containers, construction trash, and building materials were inspected to verify that the containers were tightly covered to prevent construction wastes from being dispersed by wind.

Stored Construction materials

Building materials and equipment at the DKIST site, or in lay-down or storage areas that are susceptible to wind dispersal were examined to verify that they were secured to prevent their movement by wind into native arthropod habitats.

Inspection Date

Sampling and inspection was conducted on September 7, 2017.

VI. RESULTS and DISCUSSION

Programmatic Monitoring

HIGH ALTITUDE OBSERVATORIES

The HO site covers about 18 acres and contains observatory facilities. Several areas of the site are being used to store materials and equipment. Seventy-six species of arthropods were detected at the HO site (excluding the Air Force Facility and the DKIST site). The species included seventeen endemic species, forty-five non-indigenous species, and fourteen species of unknown status.

Spiders and Mites - Arachnida

Seven species of spiders and mites were found at the HO. Two mites were uncommon in leaf litter. Juvenile and adult Lycosid spiders, *Hogna* (*Lycosa*) *hawaiiensis*, were common in pitfall traps and foraging among rocks. An unknown species of jumping spider (family Salticidae) was infrequent on rocks and cinder. Two other unknown species of spiders (families Linyphiidae and Theridiidae) were uncommon in HO sampling.

Beetles – Order Coleoptera

Sixteen beetle species were observed at the HO site, all non-indigenous. The most common beetles at the HO site were

lady-bird beetles (family Coccinellidae), including the seven-spot ladybird beetle (*Coccinella septempunctata*), the large spotted ladybird beetle (*Harmonia conformis*), the convergent ladybird beetle (*Hippodamia convergens*), and seven other less common species.



The vedalia beetle *Rodolia cardinalis* is a biocontrol agent endemic to Australia. This small ladybird beetle feeds on small arthropods such as aphids and mites.

Three species of non-indigenous leaf beetles (family Chrysomelidae) and two species of rove beetles (family Staphylinidae) were infrequent, and one species of carabid beetle (*Trechus obtusus*) was uncommon.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai'i

Flies - Order Diptera

Twelve species of flies were detected at the HO site, only two that are native to Hawai'i. An endemic species of fruit fly (*Trupanea cratericola*) was uncommon on *Dubautia* and one specimen of *Trupanea limpida* was found at Reber Circle.

The nine non-indigenous species had been collected in previous monitoring samples. The status of five species of flies collected was unknown.

True Bugs - Orders Heteroptera and Homoptera

Thirteen species of true bugs (order Heteroptera) were observed including six Hawaiian endemic species.

Endemic seed bug species *Nysius coenosulus* (family Lygaeidae) was common on both *Dubautia* and *pukiawe*. Two other endemic seed bugs (*Nysius palor* and *Nysius rubescens*) were less common. Also at the HO was an uncommon non-indigenous seed bug, *Pachybrachius fracticollis*, an introduction from Europe, where it inhabits bogs, fens and wet heathland.

Species from the family Miridae included the Hawaiian endemic insects *Engytates hawaiiensis*, abundant on *Dubautia*, *Trigonotylus hawaiiensis*, found on grasses, and *Hyalopeplus pelucidus*

Stål. A non-indigenous species of Miridae, the broken-backed bug (*Pachybrachius fracticollis*) was uncommon at the HO site.

Adults and nymphs of the non-indigenous seed bug *Geocoris pallens* were common on vegetation at the HO site. This species is a predator on small insects.



The Western bigeyed bug (*Geocoris pallens*) was common on vegetation in HO sampling.

Six species of Homoptera were found at HO, including an endemic species of plant hopper of the genus *Nesosydne*, abundant on *Dubautia*. The non-indigenous Acacia psyllid, *Acizzia uncatoides* was common. Also common was an unidentified plant hopper (family Cicadellidae).

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Bees and Wasps – Order Hymenoptera

Twelve species of bees and wasps were found at the HO site, two endemic to Hawai‘i. One endemic species of yellow-faced bees (genus *Hylaeus*) was common on *Dubautia* and *pukiawe*. Only female yellow-faced bees were collected, and because species level determination is based on male characters, identification to species was not possible.

Two specimens of a small endemic parasitoid, *Sierola* sp., were found. *Sierola* is a genus that represents a very large radiation of native parasitic wasps. The genus has been little studied, especially on Maui, where relatively few species have been described. The taxonomy of this genus is currently being revised.

Non-indigenous species found at the HO include four species of Ichneumonidae, two very small parasitoids and infrequent specimens of yellow-jackets.

Butterflies and Moths – Order Lepidoptera

Five species of Lepidoptera were found at the HO site, all endemic to Hawai‘i. These include three endemic species in the genus *Agrotis*, the common *Agrotis epicremna*, the uncommon *A. baliopa*, and the infrequent *A. mesotaxa*.

The Haleakalā flightless moth (*Thyrocopa apatela*), uncommon at the HO site, was also observed.

Other Lepidoptera included the infrequent endemic Hawaiian Blue (*Udara blackburni*).

Other Species

Two species of brown lacewings (family Hemerobiidae) were found at the HO.

Other invertebrates found at the HO site include an unknown centipede, the common woodlouse, *Porcellio scaber*, and the garlic snail (*Oxychilus alliarius*).

A complete list of arthropods observed during this sampling session at the HO site can be found in Appendix A at the end of this report. No new invasive species were observed that could impact native arthropod species. The species of indigenous arthropods detected have been observed at the site during other surveys. Conditions had been dry at the summit, and leaf litter was noticeably less moist than in previous sampling years, which seemed to correspond to lower abundance of arthropods in the soil.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai'i

DKIST CONSTRUCTION SITE

Construction was started on the DKIST in December 2012 and was ongoing during the Summer 2017 sampling session. Excavation was completed in 2014 and with one exception, earth-moving equipment has been removed from the site.

Sixty-two species of arthropods were collected at the DKIST site during the Summer 2017 sampling session. The species included sixteen endemic Hawaiian arthropods, thirty-four non-indigenous arthropods, and twelve species of unknown status.

Spiders and Mites - Arachnida

Five species of spiders and mites were found at the DKIST site. Two unknown mites were uncommon. Juvenile and adult Lycosid spiders, *Hogna hawaiiensis* Simon, were common in pitfall traps at the DKIST site, and were seen actively foraging among rocks. A small Linyphiidae spider and another spider (family Theridiidae) were uncommon.

Beetles - Order Coleoptera

Nine species of beetles were observed at the DKIST site, all non-indigenous. The species included five ladybird beetles, two leaf beetle (family Chrysomelidae), a non-indigenous Carabid beetle, and a

single specimen of the eucalyptus snout beetle (*Gonipterus scutellatus*).



The convergent ladybird beetle (*Hippodamia convergens*) was common on vegetation in HO sampling.

Flies - Order Diptera

Ten species of flies were detected at the DKIST site. Only one endemic species of fly (family Tephritidae) was observed. *Trupanea cratericola* was uncommon on pukiawe and *Dubautia*. All other species were non-indigenous in Hawai'i.

The non-indigenous species include common species such as the blue bottle fly, *Calliphora latifrons* and small flies of the family Sciaridae, and uncommon species such as the three hover flies (family Syrphidae), *Allograpta exotica*, *Eristalis tenax* and *Toxomerus marginatus*.

True Bugs - Orders Heteroptera and Homoptera

Thirteen species of true bugs (Order Heteroptera) were observed at the

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

DKIST site. These include seven Hawaiian endemic species.

Three species of the Hawaiian endemic seed bug genus *Nysius* were found at the DKIST site. *Nysius coenosulus* was common on *Dubautia* and *pukiawe*. Two other endemic species (*N. lichenicola* and *N. palor*) were infrequent in litter and on vegetation.

The common Hawaiian endemic species *Engytates hawaiiensis* (family Miridae) was observed feeding on *Dubautia* and *Trigonotylus hawaiiensis* was found on grasses. *Hyalopeplus pelucidus* is known from all major Hawaiian Islands and ranges from the seashore to several thousand feet.



Trigonotylus hawaiiensis was found on grasses at the DKIST site.

The non-indigenous species *Geocoris pallens*, was common throughout the site.

The assassin bug, *Zelus renardii*, was infrequent at the site.



The assassin bug, *Zelus renardii*.

Six species of Homoptera were collected, including a species of the endemic genus *Nesosydne* that was abundant on *Dubautia*. Non-indigenous species include a species of aphid and planthopper and the abundant Acacia psyllid.

Bees and Wasps – Order Hymenoptera

Eight species of bees and wasps were observed at the DKIST site. These species include a species of endemic yellow-faced bees, a small parasitoid, *Sierola sp.*, and an endemic wasp, *Odynerus nubicola*.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i



A small endemic parasitoid, *Sierola* sp. was present at the DKIST site.

Other Hymenoptera observed include small non-indigenous parasitic wasps.

Moths – Order Lepidoptera

Four species of Lepidoptera were collected, three endemic species and one non-indigenous species. Three large moths in the genus *Agrotis* were captured in light traps. *Agrotis epicremna* were quite abundant at both HO and DKIST, and were seen actively flying during the daytime, as well as abundant in the light traps. On one night, 62 *A. epicremna*

(and nothing else) were collected in the DKIST light trap.



The endemic *Agrotis epicremna* was very abundant.

Haleakalā flightless moths (*Thyrocopa apatela*) caterpillars were observed in leaf litter beneath vegetation.

A complete list of arthropods observed during this sampling session at the DKIST site can be found in Appendix B at the end of this report. No new invasive species were observed that could impact native arthropod species.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

HALEAKALĀ ENTRANCE STATION

Sampling in HALE occurred near the Entrance Station (HALE ES) at 6,250 feet elevation. Seventy-six species of arthropods were collected and observed there. The species included twenty-two endemic Hawaiian arthropods, forty-one non-indigenous arthropods, and thirteen species of unknown status.

Spiders and Mites - Arachnida

Two species of mites and eight species of spiders were recorded at the HALE ES site. The only species identified as endemic was a crab spider (*Mecaphesa sp. nr. kanakanus*), uncommon on vegetation.

Two jumping spiders (family Salticidae) were observed at the HALE ES. *Phidippus audax* was commonly encountered at HALE-ES, primarily as juveniles.



Phidippus audax encountered at HALE-ES.

A non-indigenous hunting spider (*Cheiracanthium mordax* L. Koch) was common in sweep net sampling over grasses. Another non-indigenous species recorded was a comb-footed spider (*Steatoda grossa* (C. L. Koch)). Similar in appearance to the black widow, this uncommon species was found under logs and rocks.

Beetles - Order Coleoptera

Nine species of beetles were observed, including an endemic ground beetle (genus *Mecyclothorax*) found under downed logs, and a similar-looking, but somewhat smaller, non-indigenous ground beetle, *Trechus obtusus*.

Non-indigenous beetles include two species of ladybird beetles and a leaf beetle (family Chrysomelidae) was observed feeding on Eucalyptus trees near the entrance station. *Paropsistrna m-fuscum* is an important pest species on Blue Gum trees.

Other non-indigenous species found were *Gonipterus scutellatus*, the gum tree weevil, native to Australia, and occurring around the World where Eucalyptus trees grow. This insect is a specialist on Eucalyptus, and it is likely that it was experiencing an outbreak on Eucalyptus at Hosmer Grove or at lower elevations. It has wings and could disperse from

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai'i

other locations. One individual was collected at DKIST on *Dubautia*, and is the first time it has been detected at the summit.



The eucalyptus snout beetle (*Gonipterus scutellatus*) on *Dubautia*.

Two other non-indigenous weevils were also observed. *Otiorhynchus cribricollis* Gyllenhal hide in cracks in the soil during the day feed on foliage at night. The Fuller rose beetle, *Pantomorus cervinus* is a leaf-eating weevil, usually feeding on leaf margins giving leaves a ragged, saw-toothed appearance.

Flies – Order Diptera

Nine species of flies were seen at the HALE ES. Seven species were from

families of common non-indigenous flies (e.g.: blow flies, and bee flies) previously reported from HALE ES. Two species were of unknown status.

True Bugs – Orders Heteroptera and Homoptera

Nine species of true bugs (Heteroptera) were found, five endemic to Hawai'i. Four endemic species from the family Miridae were collected. Three common plant bugs *Orthotylus coprosma*, on *Coprosma*, *O. sophoriodes*, on *manane*, and *Sarona* sp. were found. A fourth endemic, *Hyalopeplus pelucidus*, was uncommon, usually found only at higher elevations.

The non-indigenous *Geocoris pallens* was uncommon at the HALE ES, but was very abundant at the summit.

Six species of Homoptera were observed. Two Hawaiian endemic species of the genus *Nesosydne* (family Delphacidae), were abundant on vegetation. A third endemic species from the genus *Nesophrosyne* (family Cicadellidae), was infrequent. Non-indigenous aphids were also common on vegetation.

Bees and Wasps – Order Hymenoptera

The ten species of Hymenoptera found near the HALE Entrance Station included three infrequent endemic species, a hunting wasp (family

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

Bethylidae), a yellow-faced bee (genus *Hylaeus*), and a parasite, *Enicospilus* sp.



A female endemic *Hylaeus* bee collected at the HALE ES.

Three ants were found, *Hypoconera opaciceps*, *Linepithema humile*, and *Cardiocondyla kagutsuchi/venestula*. All have been collected during previous sampling sessions. Two non-indigenous parasites were present as well as honey bees.

Butterflies and Moths - Order Lepidoptera

Twelve species of Lepidoptera were observed at the HALE ES, nine endemic to Hawai‘i. The only common endemic was a fruit worm moth *Carposina* sp. Other endemics such as the mamane moth (*Uresiphita polygonalis virescens*)

species of *Eudonia* and *Orthomecyna*, both from the family Crambidae, and larger species of Noctuid moths were uncommon or infrequent.

The non-indigenous species were a large noctuid moth, *Pseudaletia unipunctata*, and the common lantana biocontrol, *Stenoptilodes littoralis rhynchophora*.

Other Observations

A new species to the HALE ES site was the brown lace wing *Symphorobius californicus*. This species was also found at the HO and DKIST sites and was reared from cocoons. Other arthropods were observed at the HALE ES, including centipedes, millipedes, and sowbugs common in pitfall traps, under rocks, and in decaying vegetation.

A complete list of arthropods observed during this sampling session at the HALE ES site can be found in Appendix C at the end of this report. No new invasive species were observed that could impact native arthropod species. The species of indigenous arthropods detected have been observed at the site during other surveys.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Programmatic Monitoring Discussion

The arthropods that were found during this sampling are characteristic of the fauna found during previous monitoring. No new invasive arthropods were detected at any of the sites.

No trends in populations were detected beyond normal seasonal variation and weather related abundance. The species reported are reflective only of the sites sampled, and only qualitative data of abundance were taken.

There are three Questions of Interest that are to be answered by this monitoring:

Question 1

What are the characteristic arthropod populations at the DKIST site, the larger HO site (excluding the Air Force site), and along selected areas of the HALE Road Corridor?

The Characteristic arthropods found at the monitored sites can be found in the species lists in the appendices at the end of this report.

Question 2

What adverse impacts can be detected, if any, on characteristic populations of arthropods at the DKIST site, the larger HO site (excluding the Air Force site), and along selected areas of the HALE Road Corridor that may be due to DKIST construction?

There have been minor adverse impacts on indigenous arthropod species at the monitored sites. Native vegetation was removed from the construction site during site excavation. This reduced the size of the arthropod population at the site, however, vegetation is already recovering and it can be expected that native arthropods will return to the site to exploit the renewed plant resources.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Question 3

What non-indigenous invasive arthropod species, if any, are detected at the DKIST site, the larger HO site (excluding the Air Force site), and along selected areas of the HALE Road Corridor during DKIST construction?

There were no invasive non-indigenous arthropod species detected at the HO and DKIST sites that are potentially harmful to Hawaiian native arthropods. No new invasive arthropod species were discovered at the HALE ES site.

There were three species that have not been previously collected at the HO site. They are a brown lacewing, *Symphorobius californicus*, and two parasitoids (family Ichneumonidae), *Pimpla punicipes* and *Calliephialtes grapholithae*. The brown lacewing is a purposely introduced biological control species, and the two parasitoids are adventive species. All are known from other Hawaiian Islands.

There were three species collected at the DKIST not previously recorded from that site. They are a brown lacewing, *Symphorobius californicus* (discussed above), a small leaf beetle, *Epitrix hirtipennis*, and a parasitoid, *Calliephialtes grapholithae* (discussed above). The leaf beetle occurs on all major Hawaiian Islands. Adults damage plants by chewing small rounded holes through the leaves resulting in a "shot hole" appearance. Larvae feed on roots of host plants and may tunnel into plants.

There were five species observed at the HALE Entrance Station that have not previously been found during Programmatic Monitoring. They are the brown lacewing *Symphorobius californicus* and parasitoid *Calliephialtes grapholithae* (discussed above), as well as a spider, *Neoscona theisi*, a ladybird beetle, *Diomus sp.*, and a bottle fly, *Pollenia rudis*. The spider typically builds a small orb web between branches of shrubs. While this spider is known to occur in Hawai‘i, its distribution in Hawai‘i is unknown. The ladybird beetle and bottle fly are likely an adventive species and are known from other Hawaiian Islands.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Annual Inspection

One day was allocated to the Annual Inspection during the Summer Programmatic Monitoring session. The inspection took place on September 7, 2017. During the inspection, the interiors of DKIST structures were examined for evidence of non-indigenous invasive species. No new invasive arthropod species that could potentially harm Hawaiian native arthropods were detected. The following pages describe the results of the inspection.

The following were inspected:

- DKIST Observatory Building
- DKIST Utility Building
- DKIST Site Manager’s Trailer
- DKIST Safety Officer’s Trailer
- Subcontractor Office 1
- Subcontractor Office 2
- Reber Circle Storage Containers
- Material Storage and Lay-Down Areas

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Observatory Buildings

The Observatory, attached support building, and Utility Building were inspected and found to be free of invasive, non-indigenous arthropods. Trash cans used for food waste were lined containers and had proper fitting lids. Trash cans are emptied at least once a week as needed.



The Observatory, attached support building, and Utility Building were free of non-indigenous arthropods.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i



Rooms inside the Observatory were neat and kept clean. They are swept at least once a day as needed.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i



Corners and other likely places where non-indigenous arthropods may reside was carefully inspected. There was no evidence of invasive, non-indigenous arthropods.



Trash containers in the Observatory for discarded food were lined and had properly fitting lids. Plastic bags line smaller trash cans where food may be consumed and are removed daily.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i



The exterior of the Observatory was neat, with no trash or wind-blown debris.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i



The Utility building was inspected and found to have no signs of invasive, non-indigenous arthropods.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Office Trailers

Three office trailers were inspected. The offices contain desks, chairs, and file cabinets. The trailer offices were all found to be free of non-indigenous invasive arthropod species. Trash cans in offices had lined containers used for food waste and had proper fitting lids.



The Construction Site Manager Office trailer was found to be free of non-indigenous invasive arthropods.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i



Safety Officer's Office was organized, had a lined and covered trash can that is emptied as needed. There was no evidence of any invasive arthropods.



Subcontractor Offices (1 and 2) were clean and no evidence of invasive arthropods were found.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Storage Containers

There were storage containers on the DKIST construction site, at Reber Circle and in the lower lay-down area. All were locked and not available for interior inspection.



Storage containers at the lower lay-down area were locked and not available for inspection.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Laydown and Construction Material Storage Areas

Laydown and construction material storage areas were inspected and found to be free of non-indigenous invasive arthropod species that could potentially harm Hawaiian native species. The areas are kept free of trash and wind-blown debris and material that can become wind-blown are tied down and secured to prevent movement. Only some minor material wrapping was loose and capable of being wind-blown.



Construction material storage areas were free of non-indigenous invasive arthropods.
The areas are also free of trash and wind-blown debris.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Earth-moving Equipment

There was only one piece of earth-moving equipment left at the site. It has been at the site since the beginning of construction. It is clean and has no visible signs of invasive, non-indigenous arthropods.

Outdoor Construction Waste Containers

There are two large drop-off trash containers at the construction site. Both had secure lids and are dumped when full. The containers were inspected and found to be free of non-indigenous invasive arthropod species.



Temporary trash container adjacent to the Observatory building. The container is used for temporary daily waste storage and is dumped into the larger drop-off containers daily.

Sanitary Control

Food waste is disposed of in special lined trash cans with tight fitting lids. No outdoor food waste containers were found.

ANNUAL INSPECTION DISCUSSION



The DKIST site is kept clean and neat.

The DKIST construction site and surrounding lay-down/storage areas were found to be well organized and kept neat and clean. There were no signs of invasive non-indigenous arthropods within the HO and DKIST sites that could potentially harm Hawaiian native species.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

There are three Questions of Interest that are to be answered by this annual inspection.

Question 1

What non-indigenous arthropod species, if any, are detected within the interior of DKIST facilities and the grounds within 100 ft. (30 m) of the buildings?

There were no invasive, non-indigenous arthropods found in or around the DKIST buildings that were potentially threatening to native flora and fauna.

Question 2

What non-indigenous invasive arthropod species, if any, are detected at the DKIST construction, lay-down and staging areas?

There were no invasive, non-indigenous arthropods found within the DKIST construction site, lay-down and staging areas that were potentially threatening to native flora and fauna.

Question 3

Are mitigation measures implemented that prevent the establishment of invasive species due to DKIST construction activities?

All mitigation measures were found to be implemented to prevent establishment of invasive species due to construction activities.

Earthmoving equipment were free of large deposits of soil, dirt and vegetation debris that could harbor alien arthropods. All construction materials, crates, shipping containers, packaging material, and observatory equipment were free of alien arthropods. Sanitary control of food and garbage is preventing use by invading ants and yellow jackets. Trash receptacles for food waste have attached lids and plastic liners. Their contents are collected frequently to reduce food availability for alien predators. The roll-off containers have heavy, hinged lids to prevent wind dispersal of garbage. Refuse is collected on a regular basis to ensure containers do not become full or overflow.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
and Haleakalā National Park and Annual Inspection of the DKIST
Facilities and Grounds, Maui, Hawai‘i

Trash containers inside the Observatory and Utility buildings used for food waste were clean and lined with plastic bags. Plastic bag liners are used in all garbage containers receiving food to contain leaking fluids.



Construction materials stored at the site that are susceptible to becoming wind-blown were covered with tarps, or anchored in place. There were a few small pieces of wind-blown trash in the habitat surrounding the site. Some tarps were loose and held down with pieces of lumber.

As required by the HO Management Plan, large trucks, tractors, and other heavy equipment and construction materials were inspected before entering the Park. Inspections were conducted by HALE personnel in coordination with the DKIST Construction Administrator. A log of the inspections are maintained by HALE and was not reviewed for this inspection. A log of DKIST inspection requests is maintained by the DKIST Construction Administrator at DKIST offices in Pukalani.

The Construction Site Manager and his crew do an excellent job ensuring that the mitigation measures are implemented, that the construction site is a safe place to work, and DKIST construction has a minimal impact on the surrounding habitat.

VII. LITERATURE CITED

- Ashlock, P.D. 1966. New Hawaiian Orsillinae (Hemiptera-Heteroptera: Lygaeidae). Pacific Insects 8(4): 805-825.
- Asquith, A. 1994. Revision of the endemic Hawaiian genus *Sarona* Kirkaldy (Heteroptera: Miridae: Orthotylinae). Bishop Mus. Occas. Pap. 40:1-81.
- Asquith, A. 1997. Hawaiian Miridae (Hemiptera: Heteroptera): The Evolution of Bugs and Thought. Pacific Science 51 (4): 356-365
- Beardsley, J.W. 1961. A review of the Hawaiian Braconidae (Hymenoptera). Proceedings of the Hawaiian Entomological Society 17(3): 333-366.
- Beardsley, J.W. 1966. Investigations of *Nysius* spp. And other insects at Haleakalā, Maui during 1964 and 1965. Proceedings of the Hawaiian Entomological Society 19(2):187-200.
- Beardsley, J.W. 1969. The Anagryina of the Hawaiian Islands (Hymenoptera: Encyrtidae) with descriptions of two new species. Proceedings of the Hawaiian Entomological Society 20(2): 287-310.
- Beardsley, J.W. 1976. A synopsis of the Encyrtidae of the Hawaiian Islands with keys to genera and species (Hymenoptera: Chalcidoidea). Proceedings of the Hawaiian Entomological Society 22(2): 181-228.
- Beardsley, J.W. 1977. The *Nysius* Seed Bugs of Haleakalā National Park, Maui (Hemiptera: Lygaeidae: Orsillinae). Proceedings of the Hawaiian Entomological Society 22:443-450.
- Borror, D.J., C.A. Triplehorn, and N.F. Johnson. An Introduction to the Study of Insects. Sixth Edition. Saunders College Press, San Francisco.
- Christiansen, K. and P. Bellinger. 1992. Insects of Hawai‘i Collembola. Volume 15. University of Hawai‘i Press, Honolulu. 445 pp.
- Cole, F. R., A. C. Medeiros, L. L. Loope, and W. W. Zuehlke. 1992. Effects of the Argentine ant on arthropod fauna of Hawaiian high-elevation shrubland. Ecology 73(4): 1313-1322

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai'i

- Cushman, R.A. 1944. The Hawaiian species of *Enicospilus* and *Abanchogastra* (Hymenoptera: Ichneumonidae). *Proc Haw Ent Soc* 12(1): 39-56.
- Daly, H.V. and K.N. Magnacca 2003 Hawaiian *Hylaeus* (*Nesoprosopis*) Bees (Hymenoptera: Apoidea) Volume 17. University of Hawai'i Press, Honolulu. 234 pp.
- Fullaway, D.T. & N.L.H. Krauss. 1945. Common Insects of Hawai'i. Tongg Publishing Co., Honolulu. 228 pp.
- Gagne, W.C. 1997. Insular Evolution, Speciation, and Revision of the Hawaiian Genus *Nesiomiris* (Hemiptera: Miridae). Bishop Museum Bulletin in Entomology 7. Bishop Museum Press, Honolulu.
- Gambino, P. 1992. Yellowjacket (*Vespula pensylvanica*) Predation at Hawai'i Volcanoes and Haleakala National Parks: Identity of Prey Items. *Proceedings, Hawaiian Entomological Society* Vol 31:157-164.
- Giambelluca, T.W., M.A. Nullet and T.A. Schroeder. 1986. Rainfall atlas of Hawai'i. Report R76. Hawai'i Department of Land and Natural Resources, Division of Water and Land Development, Honolulu.
- Hardy, D.E. 1960. Diptera: Nematocera-Brachycera. Insects of Hawai'i. A manual of the insects of the Hawaiian Islands, including an enumeration of the species and notes on their origin, distribution, hosts, parasites, etc. Volume 10. Diptera: Nematocera--Brachycera. University of Hawai'i Press, Honolulu. ix + 368 pp.
- Hardy, D.E. 1965. Diptera: Cyclorrhapha II, series Schizophora, section Acalypterae I, family Drosophilidae. Insects of Hawai'i. A manual of the insects of the Hawaiian Islands, including an enumeration of the species and notes on their origin, distribution, hosts, parasites, etc. Volume 12. University of Hawai'i Press, Honolulu. vii + 814 pp.
- Hardy, D.E. 1966. Descriptions and notes on Hawaiian Drosophilidae (Diptera). Pp. 195-244 In: M.R. Wheeler (ed.). *Studies in genetics. III. Morgan centennial issue.* The University of Texas, Austin. vi + 563 pp.
- Hardy, D.E. 1981. Diptera: Cyclorrhapha IV, series Schizophora, section Calyptratae. Insects of Hawai'i. A manual of the insects of the Hawaiian Islands, including an enumeration of the species and notes on their origin, distribution, hosts, parasites, etc. Volume 14. University of Hawai'i Press, Honolulu. vi + 491 pp.
- Hardy, D.E. & M.D. Delfinado. 1980. Diptera: Cyclorrhapha III, series Schizophora, section Acalypterae, exclusive of family Drosophilidae. Pp. 1-451 In: *Insects of Hawai'i.*

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

- Polhemus, D. A. 2003. An initial review of *Orthotylus* in the Hawaiian Islands, with descriptions of twenty-two new species (Heteroptera: Miridae). Journal of the New York Entomological Society 110(3-4): 270-340.
- Polhemus, D. A. 2005. Further studies on the genus *Orthotylus* (Heteroptera: Miridae) in the Hawaiian Islands, with descriptions of thirty-four new species. Journal of the New York Entomological Society 112(4): 227-333.
- Polhemus, D. A. 2011. Continuing studies on the genus *Orthotylus* in the Hawaiian Islands (Heteroptera: Miridae), with descriptions of thirty-two new species. Entomologica Americana 117(1/2): 37-109.
- Polhemus, D. A. 2013. A Fourth Contribution to the Study Of Hawaiian *Orthotylus* (Heteroptera: Miridae): New Species, Addenda, Revised Checklist, And Key To The *Perrottetia*-Feeding Species. Entomologica Americana 119(1-2):30-41.
- Sharp (ed). 1899-1913. Fauna Hawaiiensis. Cambridge-at-the-University-Press.
- Tentorio, J.M. 1969. Insects of Hawai‘i Volume 11, Supplement. Diptera: Dolichopodidae Appendix (Phoridae). University of Hawai‘i Press, Honolulu. 73 pp.
- Townes, H. 1958. Insects of Micronesia Hymenoptera: Ichneumonidae, Stephanidae, and Evaniidae. Insects of Micronesia 19(2):35-87. B.P. Bishop Museum, Honolulu.
- USFWS. 2011. Biological Opinion Of The U.S. Fish And Wildlife Service For Construction And Operation Of The Advanced Technology Solar Telescope (ATST) At The Haleakala High Altitude Observatory Site Maui, Hawai‘i June 15, 2011 (1-2-2011-F-0085)
- Wagner, W.L., D.R. Herbst, and S.H. Sohmer. 1990. Manual of the Flowering Plants of Hawai‘i. University of Hawai‘i Press, Honolulu.
- Watanabe, C. 1958. Insects of Micronesia Hymenoptera: Eucharidae. Insects of Micronesia 19(2):1-34. B.P. Bishop Museum, Honolulu.
- Williams, F.X. 1931. Handbook of the insects and other invertebrates of Hawaiian sugar cane fields. Hawaiian Sugar Planters’ Association, Honolulu. 400 pp.
- Yoshimoto, C.M. 1965a. Synopsis of Hawaiian Eulophidae including Aphelininae (Hym.: Chalcidoidea). Pac Ins 7(4): 665-699.

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

APPENDIX A

HO ARTHROPOD SPECIES LIST

A list of Arthropod species detected during the Summer 2017 sampling at the HO site.

Class	Order	Family	Genus	Species	Authority	Status
Arachnida	Acari		SP1			unknown
Arachnida	Acari		SP2			unknown
Arachnida	Araneae	Linyphiidae				unknown
Arachnida	Araneae	Lycosidae	Hogna	hawaiiensis	Simon	endemic
Arachnida	Araneae	Salticidae				unknown
Arachnida	Araneae	Theridiidae				unknown
CHILOPODA	Lithobiomorpha					unknown
Crustacea	Isopoda	Porcellionidae	Porcellio	scaber	Latreille	non-indigenous
Gastropoda	Stylommatophora	Zonitidae	Oxychilus	allarius	(J.S. Miller)	non-indigenous
Insecta	Coleoptera	Carabidae	Trechus	obtus	Erichson	non-indigenous
Insecta	Coleoptera	Chrysomelidae	Altica	carinata	(Germar)	non-indigenous
Insecta	Coleoptera	Chrysomelidae	Diachus	auratus	(Fabricius)	non-indigenous
Insecta	Coleoptera	Chrysomelidae	Paropsisterna	m-fuscum		non-indigenous
Insecta	Coleoptera	Coccinellidae	Coccinella	californica	(Mannerheim)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Coccinella	septempunctata	Linnaeus	non-indigenous
Insecta	Coleoptera	Coccinellidae	Diomus	notescens	(Blackburn)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Harmonia	conformis	(Boisduval)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Hippodemia	convergens	Gurein-Meneville	non-indigenous
Insecta	Coleoptera	Coccinellidae	Hyperaspis	pantherina or sylvestrii		non-indigenous
Insecta	Coleoptera	Coccinellidae	Olla	v-nigrum	(Mulsant)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Rhyzobius	forestieri	(Mulsant)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Rodolia	cardinalis	(Mulsant)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Scymnus	loewii	Mulsant	non-indigenous
Insecta	Coleoptera	Staphylinidae	Philonthus	sp.		non-indigenous
Insecta	Coleoptera	Staphylinidae				unknown
Insecta	Diptera	Calliphoridae	Calliphora	latifrons	Hough	non-indigenous
Insecta	Diptera	Calliphoridae	Calliphora	vomitaria	(Linnaeus)	non-indigenous
Insecta	Diptera	Chamaemyiidae	Leucopis	albipuncta	Zetterstedt	non-indigenous
Insecta	Diptera	Drosophilidae	Drosophila	melanogaster	Meigen	non-indigenous
Insecta	Diptera	Sciaridae				unknown

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

Class	Order	Family	Genus	Species	Authority	Status
Insecta	Diptera	Sepsidae	Sepsis	thoracica	(Robineau-Desvoidy)	non-indigenous
Insecta	Diptera	Syrphidae	Allograpta	exotica	(Weidemann)	non-indigenous
Insecta	Diptera	Syrphidae	Eristalis	tenax	(Linnaeus)	non-indigenous
Insecta	Diptera	Syrphidae	Toxomerus	marginatus	(Say)	non-indigenous
Insecta	Diptera	Tachinidae	Chaetogaedia	monticola	(Bigot)	non-indigenous
Insecta	Diptera	Tephritidae	Trupanea	cratericola	(Grimshaw)	endemic
Insecta	Diptera	Tephritidae	Trupanea	limpidapex	(Grimshaw)	endemic
Insecta	Heteroptera	Anthocoridae				unknown
Insecta	Heteroptera	Geocoridae	Geocoris	pallens	Stål	non-indigenous
Insecta	Heteroptera	Lygaeidae	Nysius	coenosulus	Stål	endemic
Insecta	Heteroptera	Lygaeidae	Nysius	palor	Ashlock	endemic
Insecta	Heteroptera	Lygaeidae	Nysius	rubescens	White	endemic
Insecta	Heteroptera	Lygaeidae	Pachybrachius	nr. fracticollis		non-indigenous
Insecta	Heteroptera	Miridae	Engytatus	hawaiiensis	(Kirkaldy)	endemic
Insecta	Heteroptera	Miridae	Hyalopeplus	pelucidus	Stål	endemic
Insecta	Heteroptera	Miridae	Taylorilygus	apicalis	(Fieber)	non-indigenous
Insecta	Heteroptera	Miridae	Trigonotylus	hawaiiensis	(Kirkaldy)	endemic
Insecta	Heteroptera	Nabidae	Nabis	capsiformis	Germar	non-indigenous
Insecta	Heteroptera	Reduviidae	Zelus	renardii	Kolenati	non-indigenous
Insecta	Heteroptera	Rhyparochromidae	Brentiscerus	putoni (= australis)	(White)	non-indigenous
Insecta	Homoptera	Aphididae	SP1			non-indigenous
Insecta	Homoptera	Cercopidae	Clastoptera	xanthocephala	Germar	non-indigenous
Insecta	Homoptera	Cicadellidae	SP1			unknown
Insecta	Homoptera	Delphacidae	Nesosydne	sp.		endemic
Insecta	Homoptera	Pseudococcidae	SP 1			unknown
Insecta	Homoptera	Psyllidae	Acizzia	uncatoides	(Ferris & Klyver)	non-indigenous
Insecta	Hymenoptera	Apidae	Apis	mellifera	Linnaeus	non-indigenous
Insecta	Hymenoptera	Bethylidae	Sierola	spp.		endemic
Insecta	Hymenoptera	Braconidae	Meteorus	laphygmae	Viereck	non-indigenous
Insecta	Hymenoptera	Chalcidae	Brachymeria	obscurata	(Walker)	non-indigenous
Insecta	Hymenoptera	Colletidae	Hylaeus	sp.		endemic
Insecta	Hymenoptera	Eulophidae				unknown
Insecta	Hymenoptera	Ichneumonidae	Barichneumon	californicus	(Ashmead)	non-indigenous
Insecta	Hymenoptera	Ichneumonidae	Calliephialtes	grapholithae	(Cresson)	non-indigenous

Class	Order	Family	Genus	Species	Authority	Status
Insecta	Hymenoptera	Ichneumonidae	Gelis	tenellus	(Say)	non-indigenous
Insecta	Hymenoptera	Ichneumonidae	Pimpla	punicipes	Cresson	non-indigenous
Insecta	Hymenoptera	Ichneumonidae	Spilichneumon	superbus	(Provancher)	non-indigenous
Insecta	Hymenoptera	Vespidae	Vespula	pensylvanica	(Saussure)	non-indigenous
Insecta	Lepidoptera	Lycaenidae	Udara	blackburni	(Tuely)	endemic
Insecta	Lepidoptera	Noctuidae	Agrotis	baliope	Meyrick	endemic
Insecta	Lepidoptera	Noctuidae	Agrotis	epicremna	Meyrick	endemic
Insecta	Lepidoptera	Noctuidae	Agrotis	giffardi (or mesotoxa)		endemic
Insecta	Lepidoptera	Xyloryctidae	Thyrocopa	apatela	(Walsingham)	endemic
Insecta	Neuroptera	Hemerobiidae	SP1			unknown
Insecta	Neuroptera	Hemerobiidae	Sympherobius	californicus	Banks	non-indigenous
Insecta	Psocoptera					unknown

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

APPENDIX B

DKIST ARTHROPOD SPECIES LIST

A list of Arthropod species detected during the Summer 2017 sampling at the DKIST site.

Class	Order	Family	Genus	Species	Authority	Status
Arachnida	Acari		SP1			unknown
Arachnida	Acari		SP2			unknown
Arachnida	Araneae	Linyphiidae				unknown
Arachnida	Araneae	Lycosidae	Hogna	hawaiiensis	Simon	endemic
Arachnida	Araneae	Theridiidae				unknown
CHILOPODA	Lithobiomorpha					unknown
Crustacea	Isopoda	Porcellionidae	Porcellio	scaber	Latreille	non-indigenous
Gastropoda	Stylommatophora	Zonitidae	Oxychilus	alliarius	(J.S. Miller)	non-indigenous
Insecta	Coleoptera	Carabidae	Trechus	obtusius	Erichson	non-indigenous
Insecta	Coleoptera	Chrysomelidae	Altica	carinata	(Germar)	non-indigenous
Insecta	Coleoptera	Chrysomelidae	Epitrix	hirtipennis	(Melsheimer)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Coccinella	californica	(Mannerheim)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Coccinella	septempunctata	Linnaeus	non-indigenous
Insecta	Coleoptera	Coccinellidae	Harmonia	conformis	(Boisduval)	non-indigenous
Insecta	Coleoptera	Coccinellidae	Hippodamia	convergens	Gurein-Meneville	non-indigenous
Insecta	Coleoptera	Coccinellidae	Olla	v-nigrum	(Mulsant)	non-indigenous
Insecta	Coleoptera	Curculionidae	Gonipterus	scutellatus		non-indigenous
Insecta	Diptera	Calliphoridae	Calliphora	latifrons	Hough	non-indigenous
Insecta	Diptera	Calliphoridae	Calliphora	vomitaria	(Linnaeus)	non-indigenous
Insecta	Diptera	Chamaemyiidae	Leucopis	albipuncta	Zetterstedt	non-indigenous
Insecta	Diptera	Dolichopodidae	Chrysosoma	globiferum	(Wiedemann)	non-indigenous
Insecta	Diptera	Drosophilidae	Drosophila	melanogaster	Meigen	non-indigenous
Insecta	Diptera	Sciaridae				unknown
Insecta	Diptera	Sepsidae	Sepsis	thoracica	(Robineau-Desvoidy)	non-indigenous
Insecta	Diptera	Syrphidae	Allograpta	exotica	(Wiedemann)	non-indigenous
Insecta	Diptera	Syrphidae	Eristalis	tenax	(Linnaeus)	non-indigenous
Insecta	Diptera	Syrphidae	Toxomerus	marginatus	(Say)	non-indigenous
Insecta	Diptera	Tephritidae	Trupanea	cratericola	(Grimshaw)	endemic
Insecta	Heteroptera	Anthoridae				unknown
Insecta	Heteroptera	Geocoridae	Geocoris	pallens	Stål	non-indigenous
Insecta	Heteroptera	Lygaeidae	Nysius	coenosulus	Stål	endemic

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

Class	Order	Family	Genus	Species	Authority	Status
Insecta	Heteroptera	Lygaeidae	Nysius	lichenicola	Kirkaldy	endemic
Insecta	Heteroptera	Lygaeidae	Nysius	palor	Ashlock	endemic
Insecta	Heteroptera	Miridae	Coridromius	variegatus	(Montrouzier)	non-indigenous
Insecta	Heteroptera	Miridae	Engytatus	hawaiiensis	(Kirkaldy)	endemic
Insecta	Heteroptera	Miridae	Hyalopeplus	pelucidus	Stål	endemic
Insecta	Heteroptera	Miridae	Koanoa?			endemic?
Insecta	Heteroptera	Miridae	Taylorilygus	apicalis	(Fieber)	non-indigenous
Insecta	Heteroptera	Miridae	Trigonotylus	hawaiiensis	(Kirkaldy)	endemic
Insecta	Heteroptera	Pentatomidae	Nezara	viridula	Linnaeus	non-indigenous
Insecta	Heteroptera	Reduviidae	Zelus	renardii	Kolenati	non-indigenous
Insecta	Homoptera	Aphididae	SP1			non-indigenous
Insecta	Homoptera	Cercopidae	Clastoptera	xanthocephala	Germar	non-indigenous
Insecta	Homoptera	Cicadellidae	SP1			unknown
Insecta	Homoptera	Delphacidae	Nesosydne	sp.		endemic
Insecta	Homoptera	Pseudococcidae	SP 1			unknown
Insecta	Homoptera	Psyllidae	Acizzia	uncatoides	(Ferris & Klyver)	non-indigenous
Insecta	Hymenoptera	Apidae	Apis	mellifera	Linnaeus	non-indigenous
Insecta	Hymenoptera	Bethylidae	Sierola	spp.		endemic
Insecta	Hymenoptera	Colletidae	Hylaeus	sp.		endemic
Insecta	Hymenoptera	Eulophidae				unknown
Insecta	Hymenoptera	Ichneumonidae	Barichneumon	californicus	(Ashmead)	non-indigenous
Insecta	Hymenoptera	Ichneumonidae	Calliephialtes	grapholithae	(Cresson)	non-indigenous
Insecta	Hymenoptera	Ichneumonidae	Gelis	tenellus	(Say)	non-indigenous
Insecta	Hymenoptera	Vespididae	Odynerus	nubicola	Perkins	endemic
Insecta	Lepidoptera	Noctuidae	Agrotis	baliopa	Meyrick	endemic
Insecta	Lepidoptera	Noctuidae	Agrotis	epicremna	Meyrick	endemic
Insecta	Lepidoptera	Noctuidae	Pseudaletia	unipunctata	(Haworth)	non-indigenous
Insecta	Lepidoptera	Xyloryctidae	Thyrocopa	apatela	(Walsingham)	endemic
Insecta	Neuroptera	Hemerobiidae	SP1			unknown
Insecta	Neuroptera	Hemerobiidae	Sympherobius	californicus	Banks	non-indigenous
Insecta	Psocoptera					unknown

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

APPENDIX C

HALE ES ARTHROPOD SPECIES LIST

A list of Arthropod species detected during the Summer 2017 sampling at the
HALE Entrance Station.

Class	Order	Family	Genus	Species	Authority	Status
Arachnida	Acari		SP1			unknown
Arachnida	Acari		SP2			unknown
Arachnida	Araneae	Araneidae	Neoscona	theisi	(Walckenaer)	non-indigenous
Arachnida	Araneae	Clubionidae	Cheiracanthium	mordax	L. Koch	non-indigenous
Arachnida	Araneae	Linyphiidae				unknown
Arachnida	Araneae	Salticidae	Phidippus	audax	(Hentz)	non-indigenous
Arachnida	Araneae	Salticidae				unknown
Arachnida	Araneae	Theridiidae	Steatoda	grossa	(C. L. Koch)	non-indigenous
Arachnida	Araneae	Theridiidae				unknown
Arachnida	Araneae	Thomisidae	Mecaphesa	sp. nr. kanakanus	(Karsch)	endemic
CHILOPODA	Lithobiomorpha					unknown
Crustacea	Isopoda	Porcellionidae	Porcellio	scaber	Latreille	non-indigenous
DIPLOPODA	Julida	Cylindroiulus	latistriatus		(Curtis)	non-indigenous
Gastropoda	"Slugs"					non-indigenous
Gastropoda	Stylommatophora	Zonitidae	Oxychilus	allarius	(J.S. Miller)	non-indigenous
Insecta	Coleoptera	Carabidae	Mecyclothorax	spp.		endemic
Insecta	Coleoptera	Carabidae	Trechus	obtus	Erichson	non-indigenous
Insecta	Coleoptera	Chrysomelidae	Paropsisterna	m-fuscum		non-indigenous
Insecta	Coleoptera	Coccinellidae	Diomus	sp.		non-indigenous
Insecta	Coleoptera	Coccinellidae	Olla	v-nigrum	(Mulsant)	non-indigenous
Insecta	Coleoptera	Curculionidae	Gonipterus	scutellatus		non-indigenous
Insecta	Coleoptera	Curculionidae	Otiorhynchus	cribricollis	Gyllenhal	non-indigenous
Insecta	Coleoptera	Curculionidae	Pantomorus	cervinus	(Boheman)	non-indigenous
Insecta	Coleoptera	Staphylinidae				unknown
Insecta	Dermaptera	Forficulidae	Forficula	auricularia	Linnaeus	non-indigenous
Insecta	Diptera	Calliphoridae	Calliphora	latifrons	Hough	non-indigenous
Insecta	Diptera	Calliphoridae	Calliphora	vomitaria	(Linnaeus)	non-indigenous
Insecta	Diptera	Calliphoridae	Pollenia	rudis	(Fabricius)	non-indigenous
Insecta	Diptera	Chamaemyiidae	Leucopis	albipuncta	Zetterstedt	non-indigenous

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

Class	Order	Family	Genus	Species	Authority	Status
Insecta	Diptera	Sciaridae				unknown
Insecta	Diptera	Sepsidae	Sepsis	thoracica	(Robineau-Desvoidy)	non-indigenous
Insecta	Diptera	Syrphidae	Allograpta	exotica	(Weidemann)	non-indigenous
Insecta	Diptera	Syrphidae	Toxomerus	marginatus	(Say)	non-indigenous
Insecta	Diptera	Tipulidae	SP1			unknown
Insecta	Heteroptera	Geocoridae	Geocoris	pallens	Stål	non-indigenous
Insecta	Heteroptera	Lygaeidae	Nysius	coenosulus	Stål	endemic
Insecta	Heteroptera	Lygaeidae	Pachybrachius	nr. fracticollis		non-indigenous
Insecta	Heteroptera	Miridae	Hyalopeplus	pelucidus	Stål	endemic
Insecta	Heteroptera	Miridae	Orthotylus	coprosmaphila	Polhemus	endemic
Insecta	Heteroptera	Miridae	Orthotylus	sophoriodes	Polhemus	endemic
Insecta	Heteroptera	Miridae	Sarona	sp.		endemic
Insecta	Heteroptera	Nabidae	Nabis	capsiformis	Germar	non-indigenous
Insecta	Heteroptera	Rhyparochromidae	Brentiscerus	putoni (= australis)	(White)	non-indigenous
Insecta	Homoptera	Aphididae	SP1			non-indigenous
Insecta	Homoptera	Cicadellidae	Nesophrosyne	sp. 1		endemic
Insecta	Homoptera	Delphacidae	Nesosydne	sp.		endemic
Insecta	Homoptera	Delphacidae	Nesosydne	sp. 2		endemic
Insecta	Homoptera	Pseudococcidae	SP 1			unknown
Insecta	Homoptera	Psyllidae	Acizzia	uncatoides	(Ferris & Klyver)	non-indigenous
Insecta	Hymenoptera	Apidae	Apis	mellifera	Linnaeus	non-indigenous
Insecta	Hymenoptera	Bethylidae	Sierola	spp.		endemic
Insecta	Hymenoptera	Colletidae	Hylaeus	sp.		endemic
Insecta	Hymenoptera	Eulophidae				unknown
Insecta	Hymenoptera	Formicidae	Cardiocondyla	kagutsuchi/venestula		non-indigenous
Insecta	Hymenoptera	Formicidae	Hypoconera	opaceps	(Mayr)	non-indigenous
Insecta	Hymenoptera	Formicidae	Linepithema	humile	(Mayr)	non-indigenous
Insecta	Hymenoptera	Ichneumonidae	Calliephialtes	grapholithae	(Cresson)	non-indigenous
Insecta	Hymenoptera	Ichneumonidae	Enicospilus			endemic
Insecta	Hymenoptera	Ichneumonidae	Gelis	tenellus	(Say)	non-indigenous
Insecta	Lepidoptera	Carposinidae	Carposina	sp. A		endemic
Insecta	Lepidoptera	Crambidae	Eudonia	spp.		endemic
Insecta	Lepidoptera	Crambidae	Nomophila	noctuella	(Denis & Schiffermüller)	non-indigenous
Insecta	Lepidoptera	Crambidae	Udea	heterodoxa	(Meyrick)	endemic

Programmatic Arthropod Monitoring at the Haleakalā High Altitude Observatories
 and Haleakalā National Park and Annual Inspection of the DKIST
 Facilities and Grounds, Maui, Hawai‘i

Class	Order	Family	Genus	Species	Authority	Status
Insecta	Lepidoptera	Crambidae	Udea	pyranthes	(Meyrick)	endemic
Insecta	Lepidoptera	Crambidae	Uresiphita	polygonalis	(Butler)	endemic
Insecta	Lepidoptera	Geometridae	Eupithecia	monticolans	Butler	endemic
Insecta	Lepidoptera	Geometridae	Scotorythra	sp.		endemic
Insecta	Lepidoptera	Noctuidae	Agrotis	epicremna	Meyrick	endemic
Insecta	Lepidoptera	Noctuidae	Agrotis	xiphias	Meyrick	endemic
Insecta	Lepidoptera	Noctuidae	Pseudaletia	unipunctata	(Haworth)	non-indigenous
Insecta	Lepidoptera	Pterophoridae	Stenoptilodes	littoralis	(Meyrick)	non-indigenous
Insecta	Neuroptera	Hemerobiidae	SP1			unknown
Insecta	Neuroptera	Hemerobiidae	Symphorobius	californicus	Banks	non-indigenous
Insecta	Odonata	Aeshnidae	Anax	junius	(Drury)	non-indigenous
Insecta	Orthoptera	Gryllidae	Trigonidomorpha	sjostedti	Chopard	non-indigenous
Insecta	Psocoptera					unknown